

Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/43

Paper 4 Theory (Extended)

May/June 2023

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

[Total: 6]

1	Some sy	vmbol e	guations	and word	equations	A to J	are shown
	OUTTIC 3		qualions	and word	- cquations.	, A (O O	, are showin

- $\mathbf{A} \quad \mathsf{H}^{\scriptscriptstyle +} \, + \, \mathsf{OH}^{\scriptscriptstyle -} \, \rightarrow \, \mathsf{H}_2\mathsf{O}$
- **B** $Cr^{3+} + 3OH^{-} \rightarrow Cr(OH)_{3}$
- **C** methane + chlorine → chloromethane + hydrogen chloride
- **D** propene + bromine \rightarrow 1,2-dibromopropane
- $E \quad C_{10}H_{22} \rightarrow C_8H_{18} + C_2H_4$
- **F** chlorine + aqueous potassium bromide → bromine + aqueous potassium chloride
- **G** methane + oxygen → carbon monoxide + water
- **H** $C_2H_5COOH + CH_3OH \rightarrow C_2H_5COOCH_3 + H_2O$
- I hydrogen + oxygen → water
- **J** $6CO_2 + 6H_2O \rightarrow C_6H_{12}O_6 + 6O_2$

Use the equations to answer the questions that follow.

Each equation may be used once, more than once, or not at all.

Give the letter, **A** to **J**, for the equation which represents:

(a) photosynthesis
[1]

(b) an addition reaction
[1]

(c) a precipitation reaction
[1]

(d) incomplete combustion
[1]

(e) a displacement reaction
[1]

(f) a substitution reaction
[1]

Question 2 starts on the next page.

2 (a) The symbols of the elements in Period 3 of the Periodic Table are shown.

Na Mg Al Si P S Cl Ar

Use the symbols of the elements in Period 3 to answer the questions that follow. Each symbol may be used once, more than once, or not at all.

Give the symbol of the element that:

(i)	is present in purified bauxite	[1]
(ii)	contains atoms with a full outer shell of electrons	[1]
(iii)	is used to kill microbes in water treatment	[1]
(iv)	forms an amphoteric oxide	[1]
(v)	forms an oxide which causes acid rain	[1]
(vi)	has an oxidation number of -1 when it forms a compound with hydrogen.	
		[1]

- **(b)** The relative atomic masses of elements can be calculated from the relative masses of isotopes and their percentage abundances.
 - (i) Identify the isotope to which all relative masses are compared.

.....[1]

(ii) Table 2.1 shows the relative masses and the percentage abundances of the two isotopes in a sample of magnesium.

Table 2.1

relative mass of isotope	percentage abundance of isotope
24	85
26	15

Calculate the relative atomic mass of magnesium to **one** decimal place.

relative atomic mass = [2]

(c)	An i	ion contains 10 electrons, 13 protons and 14 neutrons.	
	(i)	State the nucleon number of the ion.	
			[1]
	(ii)	Identify the element that forms this ion.	
			[1]
		[Total:	11]

- 3 Magnesium forms ionic compounds.
 - (a) Magnesium reacts with fluorine to form the ionic compound magnesium fluoride.

 The electronic configurations of an atom of magnesium and an atom of fluorine are shown in Fig. 3.1.





Fig. 3.1

(i) lons are formed by the transfer of electrons from magnesium atoms to fluorine atoms.

Complete the dot-and-cross diagrams in Fig. 3.2 to show the electronic configurations of **one** magnesium ion and **one** fluoride ion. Show the charges on the ions.



(b) Silicon tetrachloride, $SiCl_4$, and silicon(IV) oxide, SiO_2 , are covalent compounds.

Complete the dot-and-cross diagram in Fig. 3.3 to show the electronic configuration in a molecule of silicon tetrachloride. Show outer shell electrons only.




Fig. 3.3

[2]

(c) The melting points of silicon tetrachloride and silicon(IV) oxide are shown in Table 3.1.

Table 3.1

	melting point/°C
silicon tetrachloride	-69
silicon(IV) oxide	1710

(i)	Silicon	tetrachloride	has	a low	melting	point	because	it has	weak	forces	of	attraction
	betwee	n particles.										

Name the type of particles that are held together by these weak forces of attraction.

	[1]
(ii)	Explain, in terms of structure and bonding, why silicon(IV) oxide has a high melting point.

[Total: 10]

4 Hydrogen is produced by the reaction between zinc and dilute sulfuric acid, H₂SO₄.

$$Zn(s) + H_2SO_4(aq) \rightarrow ZnSO_4(aq) + H_2(g)$$

(a) A student carries out an experiment using excess zinc and dilute sulfuric acid.

The student measures the volume of hydrogen produced at regular time intervals using the apparatus shown in Fig. 4.1.

Lumps of zinc are used.




Fig. 4.1

The rate of reaction decreases as the reaction progresses. The rate eventually becomes zero.

	(i)	Explain why the rate of reaction decreases as the reaction progresses.	
			[1]
	(ii)	Explain why the rate of reaction eventually becomes zero.	
			[1]
(b)		e experiment is repeated using powdered zinc instead of lumps of zinc. other conditions remain the same.	
	Exp	plain, in terms of collision theory, why the rate of reaction increases if powdered zinc is use	ed.
			[2]

(c) The equation for the reaction is shown.

$$Zn(s) + H_2SO_4(aq) \rightarrow ZnSO_4(aq) + H_2(g)$$

 $25.0 \,\mathrm{cm^3}$ of $2.00 \,\mathrm{mol/dm^3}$ H₂SO₄(aq) is added to excess zinc.

Calculate the volume of H_2 formed at room temperature and pressure (r.t.p.). The volume of one mole of any gas is $24 \, \text{dm}^3$ at r.t.p. Use the following steps.

• Calculate the number of moles of H₂SO₄ used.

mo		
oles of H ₂ produced.	Deduce the number of moles of H	•
mo		
l ₂ formed at r.t.p.	• Calculate the volume of H ₂ formed	•
dm ^a		
[3]		

- (d) Hydrogen can also be produced by the reaction of zinc with dilute hydrochloric acid.
 - (i) Write a symbol equation for this reaction.

.....[2]

(ii) State the test for hydrogen gas.

test

positive result[1]

[Total: 10]

This question is about electricity and chemical reactions.

5

(a)		ueous copper(II) sulfate is an electrolyte. e electrolysis of aqueous copper(II) sulfate using inert electrodes forms: copper at the cathode oxygen at the anode.	
	(i)	State what is meant by the term electrolyte.	
	(ii)	State the term given to the Roman numeral, (II), in the name copper(II) sulfate.	[1]
	(iii)	State what happens to the colour of the aqueous $copper(II)$ sulfate as this electroly progresses.	
	(iv)	Write an ionic half-equation for the formation of copper at the cathode.	[1]
			[2]
	(v)	Give the formula of the ion that forms oxygen at the anode.	[1]
(b)		e electrolysis of aqueous copper(Π) sulfate is repeated using copper electrodes.	
		ite what happens to the anode.	[1]
(c)	Spo	oons can be electroplated with silver.	
	(i)	Name the substances used as:	
		the anode (positive electrode)	
		the cathode (negative electrode)	
		the electrolyte.	 [3]
	(ii)	State two reasons why spoons are electroplated.	
		1	
		2	 [2]

(d) Hydrogen–oxygen fuel cells can be used to produce electricity to power cars. Petrol produces carbon dioxide and carbon monoxide when it powers cars.

(i)	State one adverse effect of carbon dioxide and carbon monoxide.
	carbon dioxide
	carbon monoxide
	[2]
(ii)	State one disadvantage, other than cost, of using hydrogen–oxygen fuel cells to power cars compared to using petrol.
	[1]
	[Total: 16]

6	This	question	is about	nitrogen	and co	mpounds (of nitrogen.

(a) Ammonia is manufactured by the reaction between nitrogen and hydrogen in the Haber process.

The equation is shown.

$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$

.....[1]

(ii) State the source of hydrogen for the Haber process.

______[1]

(iii) State the typical conditions used in the Haber process.

temperature°C

pressure atm

[2]

(iv) Name the catalyst used in the Haber process.

.....[1]

(v) State what is meant by the term catalyst.

(b) Ammonia is converted into nitric acid.

(i) The first stage is the conversion of ammonia into nitrogen monoxide, NO. The equation is shown.

$$4NH_3(g) + 5O_2(g) \rightleftharpoons 4NO(g) + 6H_2O(g)$$

The reaction is carried out at a temperature of 900 °C and a pressure of 7 atm. The forward reaction is exothermic.

Using explanations that do **not** involve cost:

• explain why a temperature less than 900 °C is **not** used

explain why a pressure greater than 7 atm is **not** used.

[2]

	(ii)	In the second stage, nitrogen monoxide reacts with water and oxygen to produce nitric acid
		Balance the symbol equation for the reaction.
		NO + O_2 + $H_2O \rightarrowHNO_3$ [1
(c)		tudent makes aqueous copper(II) nitrate by adding an excess of solid copper(II) carbonate filute nitric acid.
	(i)	Write the symbol equation for this reaction.
		[2
	(ii)	State two observations that indicate the copper(II) carbonate is in excess.
		1
		2
		[2
	(iii)	Name \mbox{one} compound, other than $\mbox{copper}(II)$ carbonate, that can be added to dilute nitric acid to produce aqueous $\mbox{copper}(II)$ nitrate.
		[1
		[Total: 15

_	T1 1 (1)				
/	This question	19	anout	organic	compounds
•	TINO QUOCUOTI	10	aboat	organio	compounds

(a)	Propane a	nd chlorine	react at room	temperature.	An equation	for the r	eaction is shown.
•	u,	i iopano ai		Todot at Tooli	i tomporataro.	/ III oquation		Cacacion lo ono win.

$$C_3H_8 + Cl_2 \rightarrow C_3H_7Cl + HCl$$

(i) State the condition required for this reaction.

[4]

(ii) Draw the displayed formulae of **two** structural isomers with the formula C₃H₇C*l*.

[2]

(b) Alkenes are a homologous series of hydrocarbons.

(i) State **two** characteristics that all members of the same homologous series have in common.

1

(ii) Addition polymers are made from alkenes.

Complete Fig. 7.1 to show one repeat unit of the addition polymer formed from but-2-ene.




Fig. 7.1

[2]

(c) A repeat unit of a condensation polymer is shown in Fig. 7.2. The polymer is made from two monomers.

Fig. 7.2

(i) Draw the structures of the monomers used to produce the polymer in Fig. 7.2.



2

[Total: 12]

[2]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

The Periodic Table of Elements

	=	2 He	helium 4	10	Ne	neon 20	18	Ar	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	R	radon	118	Og	oganesson
	=			6	ш	fluorine 19	17	Cl	chlorine 35.5	35	Ŗ	bromine 80	53	П	iodine 127	85	¥	astatine -	117	<u>S</u>	tennessine -
	5			8	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	<u>e</u>	tellurium 128	84	Ъ	molod –	116	^	livermorium -
	>			7	Z	nitrogen 14	15	₾	phosphorus 31	33	As	arsenic 75	51	Sb	antimony 122	83	Ξ	bismuth 209	115	Mc	moscovium -
	≥			9	ပ	carbon 12	14	S	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Pb	lead 207	114	lΗ	flerovium -
	≡			2	Ω	boron 11	13	Ρl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	<i>1</i> 1	thallium 204	113	R	nihonium —
										30	Zu	zinc 65	48	g	cadmium 112	80	Нg	mercury 201	112	ى ت	copernicium -
										59	J.	copper 64	47	Ag	silver 108	79	Αn	gold 197	111	Rg	roentgenium -
Group										28	Z	nickel 59	46	Pd	palladium 106	78	五	platinum 195	110	Ds	darmstadtium -
<u>්</u>				1						27	ပိ	cobalt 59	45	몬	rhodium 103	77	'n	iridium 192	109	¥	meitnerium -
		- エ	hydrogen 1							26	Fe	iron 56	44	Ru	ruthenium 101	92	Os	osmium 190	108	H	hassium -
							1			25	Mn	manganese 55	43	ပ	technetium -	75	Re	rhenium 186			bohrium —
				16	loqu	lass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
			Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	g	niobium 93	73	Б	tantalum 181	105	Ор	dubnium –
					atc	ē				22	i=	titanium 48	40	Zr	zirconium 91	72	Ξ	hafnium 178	104	꿆	rutherfordium -
										21	Sc	scandium 45	39	>	yttrium 89	57–71	lanthanoids		89–103	actinoids	
	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	56	Ba	barium 137	88	Ra	radium -
	_			က	=	lithium 7	11	Na	sodium 23	19	×	potassium 39	37	& S	rubidium 85	55	S	caesium 133	87	ᇁ	francium —

Lu Lu	lutetium 175	103	۲	lawrencium	I
° X				_	
e9 Tm	thulium 169	101	Md	mendelevium	1
₈₈ <u>п</u>	erbium 167	100	Fm	ferminm	I
67 Ho	holmium 165	66	Es	einsteinium	_
。 O	dysprosium 163	86	ŭ	californium	_
es Tb	terbium 159	97	BK	berkelium	_
Gd	gadolinium 157	96	Cm	curium	_
e3 Eu	europium 152	92	Am	americium	_
Sm	samarium 150	94	Pu	plutonium	_
e1 Pm	promethium -	93	δ	neptunium	_
99 P N	neodymium 144	92	\supset	uranium	238
59 P	praseodymium 141	91	Ра	protactinium	231
Ce Sg	cerium 140	06	H	thorium	232
57 La	lanthanum 139	68	Ac	actinium	1

lanthanoids

actinoids

The volume of one mole of any gas is $24\,\mathrm{dm^3}$ at room temperature and pressure (r.t.p.).